

CURRENT LISTING OF CLAIMS

1. (Original) A method for stimulating an immune response, comprising administering ex vivo to a lymphoid cell a nucleic acid molecule comprising a hematopoietic cell-specific expression element operationally linked to a nucleic acid sequence encoding one or more heterologous epitopes.
2. (Original) The method of claim 1, wherein said lymphoid cell is derived from blood or a lymphoid tissue selected from the group consisting of spleen, lymph nodes, mucosa-associated lymphoid tissue (MALT), tonsils, Payer's patches, nasal-associated lymphoid tissue (NALT), Waldeyer's ring, and urogenital lymphoid tissue.
3. (Original) The method of claim 1, wherein said expression element functions in a cell selected from the group consisting of B cell and T cell.
4. (Original) The method of claim 1, wherein said epitope stimulates an antibody response.
5. (Original) The method of claim 1, wherein said epitope stimulates a CD4 T cell response.
6. (Original) The method of claim 1, wherein said epitope stimulates a CD8 T cell response.
7. (Original) The method of claim 1, wherein said epitope stimulates a CD4 T cell response and a CD8 T cell response.
8. (Original) The method of claim 1, wherein one of said epitopes stimulates an antibody response and one or more second epitopes stimulates a CD4 T cell response and a CD8 T cell response.
9. (Original) The method of claim 1, wherein said epitope is expressed as a fusion with a cytokine.

10. (Original) The method of claim 9, wherein said cytokine is selected from the group consisting of granulocyte-macrophage colony-stimulating factor, interleukin-2, interleukin-4, interferon- $\gamma$ , interleukin-5, interleukin-6, interleukin-10 and interleukin-12.

11. (Currently amended) The method of claim 1, wherein said nucleic acid molecule encodes an immunoglobulin molecule containing said heterologous epitope, wherein said epitope is inserted within a complementarity-determining region (CDR) of said immunoglobulin molecule.

12. (Original) The method of claim 11, wherein said immunoglobulin comprises a variable region.

13. (Original) The method of claim 12, wherein said variable region is a heavy chain variable region.

14. (Original) The method of claim 12, wherein said variable region is a light chain variable region.

15. (Original) The method of claim 11, wherein said immunoglobulin molecule comprises a heavy chain.

16. (Original) The method of claim 11, wherein said immunoglobulin molecule comprises a light chain.

17. (Original) A method for stimulating an immune response, comprising administering to a lymphoid cell a nucleic acid molecule comprising a hematopoietic cell-specific expression element operationally linked to a nucleic acid sequence encoding one or more heterologous epitopes, wherein said lymphoid cell is lymphoid tissue selected from the group lymph nodes, mucosa-associated lymphoid in blood or a consisting of tissue (MALT), tonsils, Payer's patches, nasal-associated lymphoid tissue (NALT), Waldeyer's ring, and urogenital lymphoid tissue.

18. (Original) The method of claim 17, wherein said expression element functions in a cell selected from the group consisting of B cell and T cell.

19. (Original) The method of claim 17, wherein said epitope stimulates an antibody response.

20. (Original) The method of claim 17, wherein said epitope stimulates a CD4 T cell response.
21. (Original) The method of claim 17, wherein said epitope stimulates a CD8 T cell response.
22. (Original) The method of claim wherein said epitope stimulates a CD4 T cell response and a CD8 T cell response.
23. (Original) The method of claim 17, wherein one of said epitopes stimulates an antibody response and one or more second epitopes stimulates a CD4 T cell response and a CD8 T cell response.
24. (Original) The method of claim 17, wherein said epitope is expressed as a fusion with a cytokine.
25. (Original) The method of claim 24, wherein said cytokine is selected from the group consisting of granulocyte-macrophage colony- stimulating factor, interleukin-2, interleukin-4 , interferon- $\gamma$ , interleukin-5, interleukin-6, interleukin-10 and interleukin-12.
26. (Original) The method of claim 17, wherein said nucleic acid molecule encodes an immunoglobulin molecule containing said heterologous epitope, wherein said epitope is inserted within a complementarity-determining region (CDR) of said immunoglobulin molecule.
27. (Original) The method of claim 26, wherein said immunoglobulin comprises a variable region.
28. (Original) The method of claim 27, wherein said variable region is a heavy chain variable region.
29. (Original) The method of claim 27, wherein said variable region is a light chain variable region.
30. (Original) The method of claim 26, wherein said immunoglobulin molecule comprises a heavy chain.

31. (Original) The method of claim 26, wherein said immunoglobulin molecule comprises a light chain.

32. (Original) A nucleic acid molecule comprising a hematopoietic cell-specific expression element operationally linked to a nucleic acid sequence encoding a heterologous polypeptide, wherein said heterologous polypeptide comprises two or more T cell epitopes.

33. (Original) The nucleic acid of claim 32, wherein said T cell epitopes are selected from the group consisting of a CD4 and a CD8 epitope, two CD4 epitopes, and two CD8 epitopes.

34. (Original) The nucleic acid of claim 32, wherein said heterologous polypeptide further comprises one or more B cell epitopes.

35. (Original) The nucleic acid molecule of claim 32, wherein said expression element functions in a cell selected from the group consisting of B cell and T cell.

36. (Original) The nucleic acid molecule of claim 32, wherein said nucleic acid sequence encodes a polypeptide expressed as a fusion with a cytokine.

37. (Original) The nucleic acid molecule of claim 36, wherein said cytokine is selected from the group consisting of granulocyte-macrophage colony-stimulating factor, interleukin-2, interleukin-4, interferon- $\gamma$ , interleukin-5, interleukin-6, interleukin-10 and interleukin-12.

38. (Original) A nucleic acid molecule comprising a hematopoietic cell-specific expression element operationally linked to a nucleic acid sequence encoding one or more heterologous epitopes, wherein said nucleic acid sequence encodes an immunoglobulin molecule containing said one or more epitopes and wherein said one or more epitopes is inserted within a complementarity-determining region (CDR) of said immunoglobulin molecule, wherein said heterologous peptide comprises two or more T cell epitopes.

39. (Original) The nucleic acid of claim 38, wherein said T cell epitopes are selected from the group consisting of a CD4 and a CD8 epitope, two CD4 epitopes, and two CD8 epitopes.

40. (Original) The nucleic acid of claim 38, further comprising one or more B cell epitopes.

41. (Original) The nucleic acid molecule of claim 38, wherein said immunoglobulin comprises a variable region.

42. (Original) The nucleic acid molecule of claim 41, wherein said variable region is a heavy chain variable region.

43. (Currently amended) ~~the~~ The nucleic acid molecule of claim 41, wherein said variable region is a light chain variable region.

44. (Original) The nucleic acid molecule of claim 38, wherein said one or more epitopes is inserted in two CDRs.

45. (Original) The nucleic acid molecule of claim 38, wherein said epitope is expressed as a fusion with a cytokine.

46. (Original) The nucleic acid molecule of claim 45, wherein said cytokine is selected from the group consisting of granulocyte-macrophage colony-stimulating factor, interleukin-2, interleukin-4, interferon- $\gamma$ , interleukin-5, interleukin-6, interleukin-10 and interleukin-12.

47. (Original) A method of treating a condition, comprising administering a non-viral vector comprising a nucleic acid molecule comprising a B cell-specific expression element operationally linked to a nucleic acid sequence encoding a heterologous polypeptide, wherein said nucleic acid molecule is targeted to a B cell and expresses said heterologous polypeptide.

48. (Original) The method of claim 47, wherein said hematopoietic cell is targeted *ex vivo*.

49. (Original) The method of claim 47, wherein said hematopoietic cell is targeted *in vivo*.

50. (Original) The method of claim 47, wherein said heterologous polypeptide is selected from the group consisting of hormone, cytokine, clotting factor and immunoglobulin.